


Vision Diagnoses Are Common After Concussion in Adolescents

Clinical Pediatrics
1-8
© The Author(s) 2015
Reprints and permissions:
sagepub.com/journalsPermissions.nav
DOI: 10.1177/0009922815594367
cpj.sagepub.com


**Christina L. Master, MD, CAQSM^{1,2}, Mitchell Scheiman, OD³,
Michael Gallaway, OD³, Arlene Goodman, MD, CAQSM⁴,
Roni L. Robinson, RN, MSN, CRNP¹, Stephen R. Master, MD, PhD⁵,
and Matthew F. Grady, MD, CAQSM^{1,2}**

Abstract

Objective. To determine the prevalence of vision diagnoses after concussion in adolescents. **Methods.** Cross-sectional study from July 1, 2013 to February 28, 2014, of patients aged 11 to 17 years with concussion evaluated in a comprehensive concussion program. **Results.** A total of 100 adolescents were examined, with a mean age of 14.5 years. Overall, 69% had one or more of the following vision diagnoses: accommodative disorders (51%), convergence insufficiency (49%), and saccadic dysfunction (29%). In all, 46% of patients had more than one vision diagnosis. **Conclusions.** A high prevalence of vision diagnoses (accommodative, binocular convergence, and saccadic eye movement disorders) was found in this sample of adolescents with concussion, with some manifesting more than one vision diagnosis. These data indicate that a comprehensive visual examination may be helpful in the evaluation of a subset of adolescents with concussion. Academic accommodations for students with concussion returning to the classroom setting should account for these vision diagnoses.

Keywords

concussion, mild traumatic brain injury, convergence insufficiency

Background

Up to 3.6 million concussions occur annually, as estimated by the Centers for Disease Control and Prevention, and this likely represents an underestimate of the clinical problem.^{1,2} Approximately 65% of these injuries occur in the pediatric and adolescent population, 5 to 18 years of age, with the 11- to 14- and 15- to 18-year-old age groups representing the largest proportion of those injured.^{3,4} There are increasing concerns that children may be particularly vulnerable to the consequences of concussion and may have more prolonged and complicated outcomes from a cognitive and developmental perspective.⁵⁻⁹

Common problems in concussion that are observed in both the adult and pediatric populations include physical signs and symptoms (headache, dizziness, nausea, balance problems, fatigue, light and noise sensitivity, sleep problems), cognitive deficits (memory, attention, executive functioning, reaction time), and emotional issues (irritability, sadness, nervousness, anxiety and depression).^{10,11} In particular, concussion-related visual complaints, including blurred or double vision, eye fatigue, the appearance of words moving on the page,

loss of place when reading, and difficulty sustaining attention on a visual task have been reported in the adult population.¹²⁻¹⁴ While significant deficits in binocular vision (convergence), accommodative (focusing) and saccadic (eye movement) disorders have been reported in adults with concussion in both the civilian and military populations with a prevalence of up to 30% to 42%,¹⁵⁻²¹ the prevalence of visual diagnoses in adolescents with concussion is unknown. The adolescent population may be at significant risk for morbidity due to such concussion-related vision disorders because of the amount of reading and visual work involved with

¹The Children's Hospital of Philadelphia, Philadelphia, PA, USA

²Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA

³Pennsylvania College of Optometry at Salus University, Philadelphia, PA, USA

⁴St. Peter's Sports Medicine Institute, Somerset, NJ, USA

⁵Weill Cornell Medical School, New York, NY, USA

Corresponding Author:

Christina L. Master, Divisions of Orthopedics and General Pediatrics, The Children's Hospital of Philadelphia, 3400 Civic Center Boulevard, Philadelphia, PA 19104, USA.
Email: masterc@email.chop.edu

full-time schoolwork.²²⁻²⁴ A recent study using a brief Vestibular/Ocular Motor Screening Assessment (VOMS) tool that clinically assesses saccades, smooth pursuits and convergence, showed that VOMS accurately distinguished concussed patients who had symptoms of oculomotor impairment from uninjured controls, aged 9 to 18 years.²⁵ This study indicates that further investigation into the prevalence of vision diagnoses following concussion is warranted.

Reading is a complex higher order integrative function, which requires adequate accommodative, vergence, and saccadic response at the initial stage of gathering visual information. These processes are necessary to initiate and sustain visual function during reading.²⁴ Areas of the brain involved in these complex tasks include the dorsolateral prefrontal cortex (DLPC), the posterior parietal cortex, and frontal and supplemental eye fields, as well as the brainstem and cerebellum. Each of these regions can be affected by concussion injury.^{14,26} The DLPC plays a significant role in attention and working memory while also interacting with multiple other regions of the brain in carrying out tasks of executive function. In addition, from an oculomotor perspective, saccadic function and vergence are also among the many tasks influenced by the DLPC. Injury to this area may result in oculomotor deficits after concussion, which may negatively affect the task of reading, and thereby, learning in adolescents. Such deficits affecting reading have been reported in adults following concussion.¹²⁻²¹ Due to the high visual work load of adolescents in school, identifying such deficits after concussion is imperative in order to properly direct their postinjury management in school, which is their occupational setting.

The objective of this study was to use comprehensive vision testing to determine the prevalence of vision diagnoses in adolescents aged 11 to 17 years presenting to an outpatient specialty concussion program for evaluation and treatment of concussion. Secondary objectives were to determine if a symptom questionnaire, the Convergence Insufficiency Symptom Survey (CISS),^{27,28} is useful in identifying adolescent patients with vision diagnoses after concussion. The CISS was not originally designed as a screening tool for convergence insufficiency; rather, it was intended for use in clinical trials to assess change in symptoms as an outcome after treatment for convergence insufficiency. We sought to evaluate the utility of the CISS as a screening tool for visual complaints in the setting of concussion as compared with the Post Concussion Symptom Scale (PCSS), which is a validated 22-point symptom questionnaire administered independently or as part of computerized neurocognitive testing.¹¹ An additional secondary

objective was to identify any associations between visual diagnoses and changes in computerized neurocognitive testing scores after injury.

Methods

We conducted a single-center, cross-sectional study of patients who presented to the Minds Matter Concussion Program at The Children's Hospital of Philadelphia. All patients who presented for clinical care between July 1, 2013 and February 28, 2014 were eligible for the study. Our institutional review board approved this study prior to the commencement of subject enrollment and data collection. Study participants were enrolled by trained research assistants during their office visit after parental informed consent and child assent was obtained. We included patients with the medical diagnosis of concussion who were 11 to 17 years old at the time of enrollment. We excluded those who were unable to participate in vision testing for any reason, did not have corrected 20/30 visual acuity, or had a preexisting history of strabismus, amblyopia, surgery, patching, or vision therapy.

A diagnosis of concussion was based on the history of any direct or indirect force transmitted to the head resulting in the temporally-associated onset of signs or symptoms of concussion as defined by the Consensus Statement on Concussion in Sport.¹¹ Vision diagnoses were made based on objective criteria obtained in vision testing as described in Table 1.²⁹ Computerized neurocognitive testing using Immediate Post-Concussion Assessment and Cognitive Testing (ImPACT), which includes the PCSS questionnaire, was performed as part of the routine clinical evaluation of the patient for concussion.^{30,31}

Standardized historical and physical examination data,³² as well as computerized neurocognitive testing results with ImPACT, were abstracted from the electronic medical record into Research Electronic Data Capture (REDCap) for analysis. As part of the study, the Convergence Insufficiency Symptom Survey (CISS), a validated questionnaire for the assessment of change in visual symptoms after treatment for convergence insufficiency, was administered to all participants. This tool has been used as an outcome measure in several randomized clinical trials studying the treatment of convergence insufficiency. It is a 15-question instrument with Likert-type scale responses and the expected score for a normal child aged 11 to 17 years is <16.^{27,28} A standardized, comprehensive vision examination, including administration of the CISS, separate from the routine clinical assessment for concussion, was performed by either a developmental optometrist or trained research

Table 1. Diagnostic Criteria for Binocular Vision, Accommodative, and Eye Movement Diagnoses.*Convergence insufficiency*

Requires: 1

Plus at least 1 finding from 2-4

Near point of convergence of ≥ 6 cm break

Exophoria at near at least 4 pd greater than at distance

Reduced positive fusional convergence at near (< 20 pd or fails Sheard's criterion^a)Vergence facility (distance or near) ≤ 9 cpm with difficulty with base-out*Convergence excess*

Requires: 1

Plus at least 1 finding from 2-3

 ≥ 3 pd esophoria at nearReduced negative fusional convergence at near (< 8 pd or fails Sheard's criterion^a)Vergence facility (distance or near) ≤ 9 cpm with difficulty with base-in*Accommodative insufficiency*

Requires: 1 or 2

Amplitude of accommodation ≥ 2 diopters below mean for age (15-1/4 age)Monocular accommodative facility ≤ 6 cpm (difficulty with minus lenses)*Saccadic dysfunction*

Requires: 1 or 2

Developmental Eye Movement Test ratio score: 1 SD or more below the mean

Developmental Eye Movement Test error score: 1 SD or more below the mean

Abbreviations: pd, prism diopters; cpm; cycles per minute; SD, standard deviation.

^aFailing Sheard's criterion = positive fusional vergence is less than twice the near phoria.

assistants. This evaluation included an assessment of visual acuity, eye alignment (Modified Thorington Test), fusional vergence (Step Vergence Test and Vergence Facility Test), convergence (Near Point of Convergence Test), accommodative amplitude and accommodative facility (Push-up Test and +2/-2 test) and eye movements (Developmental Eye Movement Test)²⁹ (Table 1).

Descriptive statistics were calculated for the outcome measures. Fisher's exact test was used for 2-way categorical comparisons of count data. Mann-Whitney analysis for nonparametric data was performed to determine the relationship between neurocognitive test scores and vision diagnoses. All significance levels were set at $P < .05$. Statistical analysis was performed using R version 3.0.3.³² Bootstrap 95% confidence intervals (CIs) were estimated using the boot package.^{34,35} Prevalence ratios (PRs) and 95% CIs were calculated using the epiR package.³⁶

Results

One hundred participants were enrolled during the study period. The mean age was 14.5 years (age range 13.5-14.8 years). Of the study subjects, 42% were male and 65% sustained their concussion as the result of sports participation, while the remainder sustained their concussions in nonsports activities such as falls. Among the

enrolled subjects, 29% were seen within 1 month of their injury, 26% were evaluated between 1 and 3 months after their injury, and 45% were seen > 3 months after their injury. Patients seen within 1 month of injury were more likely to have a vision diagnosis than those evaluated greater than 3 months after injury. Patients who were evaluated early after injury (within 30 days) were more likely to have a vision diagnosis than those evaluated later after injury (> 90 days) ($P = .03$, PR = 1.41; 95% CI = 1.04-1.92). A total of 32% of subjects had a history of prior concussion, which was not associated with an increased risk of vision problems after concussion ($P = .36$).

The primary objective of this study was to determine the prevalence of vision diagnoses after concussion in an adolescent population age 11 to 17 years. Overall, 69% of enrolled subjects had at least one vision diagnosis after concussion, with 51% presenting with accommodative insufficiency or infacility, 49% with convergence insufficiency, and 29% with saccadic dysfunction. Of note, a substantial number of subjects had more than one vision diagnosis following concussion: 22% of patients with convergence insufficiency also had accommodation deficits, while 14% of those with both convergence insufficiency and accommodation deficits also had saccadic dysfunction on exam simultaneously (Figure 1).




Figure 1. Vision diagnoses after concussion.

Table 2. Association of Convergence Insufficiency Symptom Score (CISS) With Vision Diagnosis.

Vision Diagnosis	CISS <16	CISS ≥16
(-) Any vision diagnosis	19	12
(+) Any vision diagnosis	15	54
		<i>P</i> = .0002
(-) Convergence insufficiency	24	27
(+) Convergence insufficiency	10	39
		<i>P</i> = .006
(-) Accommodative dysfunction	23	27
(+) Accommodative dysfunction	11	39
		<i>P</i> = .02

A score of ≥ 16 on the CISS indicates that the patient may be symptomatic from convergence insufficiency. A secondary objective of this study was to determine if the CISS could be useful as a screening tool for identifying vision diagnoses in concussion as compared with the widely used PCSS. Among our study subjects, only 29% of patients specifically reported vision problems on the PCSS. In our study, the mean CISS score was 19.8 for all study participants. Overall, the mean CISS score was 22.9 for those with a vision diagnosis, as compared to a score of 13.0 in those without any vision diagnoses (Mann-Whitney $P < .001$), and a high CISS score of ≥ 16 was strongly associated with a vision diagnosis ($P < .001$, $PR = 1.85$; 95% $CI = 1.25-2.75$). (Table 2.) More specifically, a CISS score of ≥ 16 was highly correlated with the diagnosis of convergence insufficiency ($P = .006$, $PR = 2.01$; 95% $CI = 1.15-3.51$) and accommodative dysfunction ($P = .02$, $PR = 1.83$; 95% $CI = 1.08-3.09$). In contrast, a high CISS score of ≥ 16 was not associated with the presence of saccadic dysfunction ($P = .25$) (Table 2).

Another secondary objective of this study was to determine if there was any relationship between vision diagnoses after concussion and scores on computerized neurocognitive testing. Specifically, we examined the relationship between the presence of a visual diagnosis and test-derived verbal memory composite score, visual memory composite score, visual motor speed composite score, reaction time, and impulse control (Table 3). Poor verbal memory composite scores ($P = .002$) and visual motor speed scores ($P = .005$) were significantly correlated with the presence of a vision diagnosis.

Discussion

Vision problems are very common in adults following concussion,¹²⁻²¹ and vision complaints have been described in the adolescent population with concussion.²⁵ We found that a substantial proportion of adolescents in our clinical population with concussion have visual diagnoses identifiable with a comprehensive vision assessment beyond visual acuity.²⁹ Our findings indicate a much higher prevalence of vision diagnoses in adolescents with concussion, as compared with reported prevalence rates in the general pediatric population without concussion. The prevalence of convergence insufficiency in children has been reported between 2% and 8% and the prevalence of accommodative dysfunction has been reported as 5% in the general population, as compared with 49% and 50%, respectively, in our study of adolescents with concussion.³⁷⁻³⁹ While our study excluded patients with binocular vision disorders, it is possible that undiagnosed binocular vision problems comprise a portion of our patients with binocular vision problems following concussion. Despite this limitation, it is unlikely that undiagnosed preexisting binocular issues account for the notable difference observed between the prevalence of convergence insufficiency in our cohort with concussion as compared with the reported prevalence in the general population.

Oculomotor functioning is supported by the occipital lobe, brainstem, frontal lobe, parietal lobe, and cerebellum, with significant integration between these areas and the language areas of the brain to support reading.^{14,26} Because of the diffuse nature of concussion injury, both specific areas of the brain, as well as the integrative pathways needed for accurate and automatic oculomotor functioning, may be affected. The high prevalence of vision diagnoses after concussion reflects the widespread neural architecture involved in visual processing, which may make the visual oculomotor system particularly susceptible to the diffuse stretch injury seen in concussion.

Table 3. Relationship Between Neurocognitive Test Scores and Vision Diagnoses.

	(-)Vision Diagnosis	(+)Vision Diagnosis	P
Verbal memory composite score	88.4	76.9	.002
Visual memory composite score	75.8	68.0	.66
Visual motor speed composite score	37.6	32.0	.005
Reaction time	0.58	0.66	1
Impulse control	6.8	7.8	.88

Identification of concussion-related vision diagnoses is important because of the extensive visual demands of adolescents engaged in full-time school. In today's modern classroom, the use of technology is widespread and these electronic interfaces may add an additional level of visual demand when recovering from a concussion. The high prevalence of vision diagnoses following concussion in this study highlights the importance of implementing appropriate academic accommodations for students returning to the classroom.^{22,23} Since accommodative, binocular vision, and eye movement problems can cause significant symptoms related to visual activities,²⁷ it is important for physicians to suggest specific school-based accommodations to optimally manage these vision-related symptoms. Strategies that include frequent visual breaks, oral teaching, audiobooks, large-font printed material (vs. small font electronically displayed material), or preprinted notes may be extremely helpful to a student returning to learn after a concussion.^{22,23}

The PCSS is a widely used tool for diagnosis of concussion and for monitoring recovery over time.³¹ In our study population, fewer than one-third of the study subjects reported visual symptoms using this scale. In comparison, the CISS is a validated 15-point questionnaire developed as a tool to monitor changes in symptoms after treatment for convergence insufficiency.^{27,28,39} In our study, the CISS identified patients with vision diagnoses following concussion, and shows promise as a potential screening tool for vision diagnoses following concussion.

For the general pediatrician, the CISS may prove to be a highly cost-effective means of identifying patients with concussion-related vision deficits who might benefit from referral to a concussion specialist or an eye care professional for a comprehensive visual and oculomotor evaluation beyond visual acuity testing and for possible treatment of accommodation, binocular vision, and eye movements. Future studies to determine potential cutoff scores for the CISS in identifying vision diagnoses after concussion would be helpful.

Computerized neurocognitive testing with tools such as ImPACT are commonly used to aid in the diagnosis and management of acute concussions. In our study,

vision disturbances were correlated with poorer composite scores on visual motor speed, which might be expected in disorders of vision. In addition, vision diagnoses were also associated with poorer verbal memory composite scores, which may reflect the role of the DLPC in both working memory and oculomotor function. This correlation may be due to multiple factors, including the visual nature of the computerized neurocognitive test, as well as the overlapping function of the DLPC in executive function, memory, and oculomotor visual function. The fact that computerized neurocognitive testing, in general, requires sustained visual attention on a computer screen and, therefore presents a high visual demand, may prove problematic for patients with vision diagnoses following concussion. The possibility of concussion-related vision diagnoses may need to be accounted for when interpreting the ImPACT test results much in the same way that adolescents with attention or other learning disorders may have global difficulty with neurocognitive testing.⁴⁰

In the general population without concussion, there have been randomized clinical trials demonstrating the effectiveness of vergence/accommodative therapy for the treatment of convergence insufficiency and accommodative insufficiency.⁴¹⁻⁴⁴ In addition, preliminary studies have shown promise for this type of therapy for the vision deficits following concussion in adults.⁴⁵ Our results suggest that office-based vergence/accommodative therapy for vision disorders following concussion in adolescents warrants further study.

Limitations of our study include a potentially more clinically complicated sample than is seen in the general population with concussion. The Minds Matter Concussion Program at The Children's Hospital of Philadelphia is a subspecialty referral program, and as such, our cohort may represent a concussion population with a greater likelihood of prolonged/complicated recovery or vision diagnoses than is seen in the general pediatric concussion population. Although we did not have a control group, published prevalence rates of these vision diagnoses in the general pediatric population are much lower than the high prevalence of vision diagnoses following concussion in our cohort.

In our study, the 29% of patients less than 1 month postinjury were more likely to have vision deficits than the 22% of patients greater than 3 months postinjury, which was statistically significant. This may indicate that vision diagnoses are a significant component of immediate postconcussion symptomatology, and, in some cases, may recover spontaneously on their own. It is possible that vision diagnoses are present in the estimated 80% to 90% of concussion patients who recover spontaneously over the course of a few weeks,⁴⁶ and therefore, presumably, their vision problems have also spontaneously recovered. While vision diagnoses are also seen in those with prolonged recoveries in our study, further study is needed to determine if they represent a negative prognostic predictor for complicated concussion recovery. Future study is also needed to determine which symptoms are an accurate predictor of the presence of these vision diagnoses, as the PCSS does not appear to identify a majority of patients with vision diagnoses while the CISS appeared to do better. A prospective longitudinal study will be required to answer these important questions.

Given the high prevalence of vision diagnoses after concussion in the adolescent population, future studies should try to establish efficient clinical screening tools, such as the VOMS or CISS, that could be used by a pediatrician to identify the subset of patients who require further concussion care and comprehensive vision evaluation. In addition, a randomized controlled clinical trial is necessary in order to evaluate the potential benefits of possible interventions, such as vergence/accommodative therapy, for patients with vision diagnoses after concussion.

Conclusion

Vision diagnoses are prevalent in adolescents with concussion and include convergence insufficiency, accommodative disorders and saccadic dysfunction. Symptoms of these problems may include double vision, blurry vision, headache, difficulty with reading or other visual work, such as the use of a tablet, smartphone, or computer monitor in the school setting. This likely represents a significant morbidity for adolescents whose primary work is school, which is heavily visually oriented. Recognition of these deficits is essential for clinicians who care for patients with concussion and the CISS may prove to be a useful screening tool for use in the future. Identification of these vision diagnoses will help physicians design necessary academic accommodations for patients who have visual deficits and are attempting to reintegrate into school and learning while recovering from concussion.

Acknowledgments

We would like to thank Alissa Beckett, William Ensor, and Julia Lockyer for their assistance with this project.

Author Contributions

CLM made a substantial contribution to the concept and design, acquisition of data and analysis and interpretation of data, drafted the article and revised it critically for important intellectual content and approved the version to be published.

MS, MG, AG, RLR and MFG made a substantial contribution to the concept and design, acquisition of data and analysis and interpretation of data, revised the article critically for important intellectual content and approved the version to be published.

SRM made a substantial contribution to the analysis and interpretation of data, revised the article critically for important intellectual content and approved the version to be published.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This study was supported by the American Medical Society for Sports Medicine Young Investigator Grant.

References

1. Langlois JA, Rutland-Brown W, Wald MM. The epidemiology and impact of traumatic brain injury: a brief overview. *J Head Trauma Rehabil.* 2006; 21:375-378.
2. Faul M, Likang X, Wald MM, Coronado VG. *Traumatic Brain Injury in the United States: Emergency Department Visits, Hospitalizations and Deaths 2002-2006.* Atlanta, GA: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Injury Prevention and Control; 2010.
3. Centers for Disease Control and Prevention. Nonfatal traumatic brain injury from sports and recreation activities—United States 2001-2005. *MMWR Morb Mortal Wkly Rep.* 2007;56(29):733-737.
4. Gilchrist J, Thomas KE, Xu L, McGuire LC, Coronado V. Nonfatal traumatic brain injuries related to sports and recreation activities among persons aged <19 years—United States 2001-2009. *MMWR Morb Mortal Wkly Rep.* 2011;60:1337-1342.
5. Zuckerman SL, Odom M, Lee YM, Forbes J, Sills AK, Solomon G. Sport-related concussion and age: number of days to neurocognitive baseline. *Neurosurgery.* 2012;71:E558.
6. Sim A, Terryberry-Spohr L, Wilson KR. Prolonged recovery of memory functioning after mild traumatic brain injury in adolescent athletes. *J Neurosurg.* 2008;108:511-516.

7. Moser RS, Schatz P, Jordan BD. Prolonged effects of concussion in high school athletes. *Neurosurgery*. 2005;57:300-306.
8. Field M, Collins MW, Lovell MR, Maroon J. Does age play a role in recovery from sports-related concussion? A comparison of high school and collegiate athletes. *J Pediatr*. 2003;142:546-553.
9. Makdissi M, Davis G, Jordan B, Patricios J, Purcell L, Putukian M. Revisiting the modifiers: how should the evaluation and management of acute concussions differ in specific groups? *Br J Sports Med*. 2013;47:314-320.
10. Harmon KG, Drezner J, Gammons M, et al. American Medical Society for Sports Medicine position statement: concussion in sport. *Clin J Sport Med*. 2013;23:1-18.
11. McCrory P, Meeuwisse WH, Aubry M, et al. Consensus statement on concussion in sport: the 4th International Conference on Concussion in Sport held in Zurich, November 2012. *Br J Sports Med*. 2013;47:250-258.
12. Capo-Aponte JE, Urosevich TG, Temme LA, Tarbett AK, Sanghera NK. Visual dysfunctions and symptoms during the subacute stage of blast-induced mild traumatic brain injury. *Mil Med*. 2012;177:804-813.
13. Goodrich GL, Kirby J, Cockerham G, Ingalla SP, Lew HL. Visual function in patients of a polytrauma rehabilitation center: a descriptive study. *J Rehabil Res Dev*. 2007;44:929-936.
14. Ciuffreda KJ, Kapoor N, Rutner D, Suhoff IB, Han ME, Craig S. Occurrence of oculomotor dysfunctions in acquired brain injury: a retrospective analysis. *Optometry*. 2007;78:155-161.
15. Brahm KD, Wilgenburg HM, Kirby J, Ingalla S, Chang CY, Goodrich GL. Visual impairment and dysfunction in combat-injured servicemembers with traumatic brain injury. *Optom Vis Sci*. 2009;86:817-825.
16. Doble JE, Feinberg DL, Rosner MS, Rosner AJ. Identification of binocular vision dysfunction (vertical heterophoria) in traumatic brain injury patients and effects of individualized prismatic spectacle lenses in the treatment of postconcussive symptoms: a retrospective analysis. *PM R*. 2010;2:244-53.
17. Green W, Ciuffreda KJ, Thiagarajan P, Szymanowicz D, Ludlam DP, Kapoor N. Accommodation in mild traumatic brain injury. *J Rehabil Res Dev*. 2010;47:183-199.
18. Goodrich GL, Flyg HM, Kirby JE, Chang CY, Martinsen GL. Mechanisms of mTBI and visual consequences in military and veteran populations. *Optom Vis Sci*. 2013;90:105-112.
19. Stelmack JA, Frith T, Van Koeveing D, Rinne S, Stelmack TR. Visual function in patients followed at a Veterans Affairs polytrauma network site: an electronic medical record review. *Optometry*. 2009;80:419-424.
20. Suhoff IB, Kapoor N, Waxman R, Ference W. The occurrence of ocular and visual dysfunctions in an acquired brain-injured patient sample. *J Am Optom Assoc*. 1999;70:301-308.
21. Dougherty AL, MacGregor AJ, Han PP, Heltemes KJ, Galarneau MR. Visual dysfunction following blast-related traumatic brain injury from the battlefield. *Brain Inj*. 2011;25:8-13.
22. Master CL, Gioia GA, Leddy JJ, Grady MF. Importance of 'return-to-learn' in pediatric and adolescent concussion. *Pediatr Ann*. 2012;41(9):1-6.
23. Halstead ME, McAvoy K, Devore CD, Carl R, Lee M, Logan K; Council on Sports Medicine and Fitness, and Council on School Health. Returning to learning following a concussion. *Pediatrics*. 2013;132:948-957.
24. Ritty J, Solan H, Cool S. Visual and sensory-motor functioning in the classroom: a preliminary report of ergonomic demands. *J Am Optom Assoc*. 1993;64:238-244.
25. Mucha A, Collins MW, Elbin RJ, et al. A brief Vestibular-ocular Motor Screening (VOMS) assessment to evaluate concussions: preliminary findings. *Am J Sports Med*. 2014;42:2479-2486.
26. Chang A, Cohen A, Kapoor N. Top-down visual framework for optometric vision therapy for those with traumatic brain injury. *Optom Vis Perf*. 2013;1(2):48-53.
27. Borsting E, Rouse MW, Deland PN, et al. Association of symptoms and convergence and accommodative insufficiency in school-age children. *Optometry*. 2003;74:25-34.
28. Borsting EJ, Rouse MW, Mitchell GL, et al. Validity and reliability of the revised Convergence Insufficiency Symptom Survey in children aged 9-18 years. *Optom Vis Sci*. 2003;80:832-838.
29. Scheiman M, Wick B. *Clinical Management of Binocular Vision: Heterophoric, Accommodative and Eye Movement Disorders*. 4th ed. Philadelphia, PA: Lippincott; 2013.
30. Lovell M. The imPACT neuropsychological test battery. In: Echemendia RJ, ed. *Sports Neuropsychology: Assessment and Management of Traumatic Brain Injury*. New York, NY: Guilford Press; 2006:193-215.
31. Pardini D, Stump J, Lovell M, Collins M, Moritz K, Fu F. The Post-Concussion Symptom Scale (PCSS): a factor analysis. *Br J Sports Med*. 2004;38:661-662.
32. Master CL, Grady MF. Office-based management of pediatric and adolescent concussion. *Pediatr Ann*. 2012;41(9):1-6.
33. R Core Team. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing; 2013. <http://www.R-project.org/>. Published March 20, 2013. Accessed March 11, 2014.
34. Canty A, Ripley B. boot: Bootstrap R (S-Plus) Functions. R package version 1.3-9-1. <https://launchpad.net/ubuntu/+source/boot/1.3-9-1>. Published March 26, 2013. Accessed March 11, 2014.
35. Davison AC, Hinkley DV. *Bootstrap Methods and Their Applications*. Cambridge, England: Cambridge University Press; 1997.
36. Stevenson M. epiR: an R package for the analysis of epidemiological data. R package version 0.9-57. URL <http://cran.r-project.org/web/packages/svSocket/index.html>. Published March 2, 2014. Accessed March 11, 2014.
37. Scheiman M, Gallaway M, Coulter S, et al. Prevalence of vision and ocular disease conditions in a clinical pediatric population. *J Am Optom Assoc*. 1996;67:193-202.
38. Létourneau JE, Ducic S. Prevalence of convergence insufficiency among elementary school children. *Can J Optom*. 1988;50:194-197.

39. Rouse MW, Borsting E, Hyman L, et al. Frequency of convergence insufficiency among fifth and sixth graders. *Optom Vis Sci.* 1999;76:643-649.
40. Zuckerman SL, Lee YM, Odom MJ, Solomon GS, Sills AK. Baseline neurocognitive scores in athletes with attention deficit-spectrum disorders and/or learning disability. *J Neurosurg Pediatr.* 2013;12:103-109.
41. Scheiman M, Mitchell GL, Cotter S, et al. A randomized trial of the effectiveness of treatments for convergence insufficiency in children. *Arch Ophthalmol.* 2005;123:14-24.
42. Convergence Insufficiency Treatment Trial Investigator Group. A randomized clinical trial of treatments for symptomatic convergence insufficiency in children. *Arch Ophthalmol.* 2008;126:1336-1349.
43. Scheiman M, Gwiazda J, Li T. Non-surgical interventions for convergence insufficiency. *Cochrane Database Syst Rev.* 2011;(3):CD006768.
44. Scheiman M, Cotter S, Kulp MT, et al. Treatment of accommodative dysfunction in children: results from a randomized clinical trial. *Optom Vis Sci.* 2011;88:1343-1352.
45. Thiagarajan P, Ciuffreda KJ. Effect of oculomotor rehabilitation on vergence responsivity in mild traumatic brain injury. *J Rehabil Res Dev.* 2013;50:1223-1240.
46. McCrea M, Guskiewicz K, Randolph C, et al. Incidence, clinical course, and predictors of prolonged recovery time following sport-related concussion in high school and college athletes. *J Int Neuropsychol Soc.* 2013;19:22-33.